UMI

SHOULD READ 1998/1999

ELSEVIER

Biomembranes

Vol. 1415, 1999

BIOCHIMICA ET BIOPHYSICA ACTA

BBA

Author Index

Altendorf, K., see Heermann, R. (1415) 114
Altendorf, K., see Gaßel, M. (1415) 77
Auger, M., see Bouchard, M. (1415) 181

Benning, C., see Härtel, H. (1415) 205
Bouchard, M., Le Guernevé, C. and Auger, M.
Comparison between the dynamics of lipid/gramicidin A systems in the lamellar and hexagonal phases: a solid-state ^{13}C NMR study (1415) 181
Burson, K.K., see Hall, M.P. (1415) 101

Cabral-Lilly, D., see Li, X. (1415) 23
Cailler, F., Howell, S. and Crine, P.
Role of the glycosyl-phosphatidylinositol anchor in the intracellular transport of a transmembrane protein in Madin-Darby canine kidney cells (1415) 1
Cajal, Y., see Oh, J.-T. (1415) 235
Chan, H.C., see So, S.C. (1415) 261
Cheng, C.Y., see So, S.C. (1415) 261
Cheng, Y., see Hiroaki, Y. (1415) 361
Chung, Y.W., see So, S.C. (1415) 261
Crine, P., see Cailler, F. (1415) 1
Cruz, A., Marsh, D. and Pérez-Gil, J.
Rotational dynamics of spin-labelled surfactant-associated proteins SP-B and SP-C in dipalmitoylphosphatidylcholine and dipalmitoylphosphatidylglycerol bilayers (1415) 125

de Kruijff, B., see Harris, F. (1415) 10
Demel, R., see Harris, F. (1415) 10
Deslauriers, R., see Gruwel, M.L.H. (1415) 41
Dhurjati, P.S., see Oh, J.-T. (1415) 235
Duclohier, H., Snook, C.F. and Wallace, B.A.
Antiamoebin can function as a carrier or as a pore-forming peptaibol (1415) 255

Eckstein-Ludwig, U., see Liu, Y. (1415) 246
Epstein, W., see Gaßel, M. (1415) 77
Essigmann, B., see Härtel, H. (1415) 205

Fei, J., see Liu, Y. (1415) 246
Franzin, C.M., Macdonald, P.M., Polozova, A. and Winnik, F.M.
Destabilization of cationic lipid vesicles by an anionic hydrophobically modified poly(*N*-isopropylacrylamide) copolymer: a solid-state ^{31}P NMR and ^{2}H NMR study (1415) 219
Fromm, M.F., Leake, B., Roden, D.M., Wilkinson, G.R. and Kim, R.B.
Human MRP3 transporter: identification of the 5′-flanking region, genomic organization and alternative splice variants (1415) 369
Fujiyoshi, Y., see Hiroaki, Y. (1415) 361

Gaßel, M., Siebers, A., Epstein, W. and Altendorf, K.
Assembly of the Kdp complex, the multi-subunit K^+-transport ATPase of *Escherichia coli* (1415) 77
Gioioso, S., see Moore, D.J. (1415) 342
Grima, J., see So, S.C. (1415) 261
Gruner, S.M., see Li, X. (1415) 23
Gruwel, M.L.H., Kuzio, B., Xiang, B., Deslauriers, R. and Kupriyanov, V.V.
Temperature dependence of monovalent cation fluxes in isolated rat hearts: a magnetic resonance study (1415) 41

Hall, M.P., Burson, K.K. and Huestis, W.H.
Interactions of a vesicular stomatitis virus G protein fragment with phosphatidylserine: NMR and fluorescence studies (1415) 101
Harauz, G., see Palaniyar, N. (1415) 85
Harris, F., Demel, R., de Kruijff, B. and Phoenix, D.A.
An investigation into the lipid interactions of peptides corresponding to the C-terminal anchoring domains of *Escherichia coli* penicillin-binding proteins 4, 5 and 6 (1415) 10
Härtel, H., Essigmann, B., Lokstein, H., Hoffmann-Benning, S., Peters-Kottig, M. and Benning, C.
The phospholipid-deficient *pho1* mutant of *Arabidopsis thaliana* is affected in the organization, but not in the light acclimation, of the thylakoid membrane (1415) 205
Heermann, R., Altendorf, K. and Jung, K.
The turgor sensor KdpD of *Escherichia coli* is a homodimer (1415) 114
Heimburg, T.
Mechanical aspects of membrane thermodynamics. Estimation of the mechanical properties of lipid membranes close to the chain melting transition from calorimetry (1415) 147
Hernández-Hernández, A., see Rocío Marcos, M. (1415) 163
Hiroaki, Y., Mitsuoka, K., Cheng, Y., Hiroaki, H. and Fujiyoshi, Y.
Influence of various nucleotides on the in situ crystallization of Ca^{2+}-ATPase (1415) 361
Hiroaki, H., see Hiroaki, Y. (1415) 361
Hirsh, D.J., see Li, X. (1415) 23
Hoffmann-Benning, S., see Härtel, H. (1415) 205
Howell, S., see Cailler, F. (1415) 1
Huang, S.-G., see Klingenberg, M. (1415) 271
Huestis, W.H., see Hall, M.P. (1415) 101
Hui, S.W., see Langner, M. (1415) 323
Hunter, G.W. and Squier, T.C.
Phospholipid acyl chain rotational dynamics are independent of headgroup structure in unilamellar vesicles containing binary mixtures of dioleoyl-phosphatidylcholine and dioleoyl-phosphatidylethanolamine (1415) 63

Ikigai, H., Ono, T., Nakae, T., Otsuru, H. and Shimamura, T.
Two forms of *Vibrio cholerae* O1 El Tor hemolysin derived from identical precursor protein (1415) 297
Ivanov, I.T.
Low pH-induced hemolysis of erythrocytes is related to the entry of the acid into cytosole and oxidative stress on cellular membranes (1415) 349